


Clark's Grebe *Aechmophorus clarkii*

		Summer range map	Winter range map
 <p>Photo by Peter S. Weber</p>	Not Available	Not Available	Not Available

Migration Status	Breeding Habitat	Nest Location	Nest Type	Clutch Size	Incubation	Fledge	Number of Broods	Diet
N/A	Wetland-open water	N/A	N/A	3-4	23 days	63-77	1	Aquatic Invertebrates, Fish

Identification Tips: <ul style="list-style-type: none"> ● Length: 18? inches Wingspan: 40? inches ● Sexes similar ● A large, long-necked grebe with a long bill ● Bill yellow ● Red eye ● Black crown and nape ● White chin, throat, face, and belly ● White face includes eye ● Dark blackish-brown back and wings ● White secondaries ● Feet set far back on body 				Similar species: Loons are larger with longer bodies and larger, darker bills. Horned and Eared Grebes are smaller with much shorter bills that are never yellow. Red-necked Grebe is smaller without the bold black and white neck pattern. The formerly conspecific Westerns Grebe is very similar but has a greener bill. The dark crown of the Western Grebe includes the eye, whereas in the Clark's it does not. The two species have distinctive calls: Clark's give a single "kreeet" while the Western gives a doubled note. Hybridization is known go atypical birds should be studied well for exact bill color and face pattern.				
				Credits: Citation: Gough, G.A., Sauer, J.R., Iliff, M. <i>Patuxent Bird Identification Infocenter</i> 1998. Version 97.1. Patuxent Wildlife Research Center, Laurel, MD. http://www.mbr-pwrc.usgs.gov/Infocenter/infocenter.html Length and wingspan from: Robbins, C.S., Bruun, B., Zim, H.S., (1966). <i>Birds of North America</i> . New York: Western Publishing Company, Inc. Clutch size, fledging, brood, and incubation information from: Ehrlich, P., Dobkin, D., and Wheye, D. (1988). <i>The Birder's Handbook: A Field Guide to the Natural History of North American Birds</i> . New York. Simon and Schuster Inc.				